Acommendations on Auxiliary Acouncture Treatment of COVID-19

American Association of Chinese Medicine and Acupuncture has formulated 2020.3.31

Disclaimer

These recommendations are intended for consideration by licensed Acupuncturists, and health care providers who are properly trained and legally certified in acupuncture, when choosing acupoints for alleviating some common symptoms caused by COVID-19.

There are no FDA-approved therapeutics or drugs to treat, cure or prevent COVID-19. These recommendations only suggest that acupuncture may help alleviate some common symptoms of the disease from a supportive care perspective.

These recommendations are not intended to replace your doctor's diagnosis and care, or to supersede any of your doctor's advice or prescriptions.

You are responsible for consulting your doctor before trying any new therapeutics.

Contents

Recommendation on Ear Acupuncture



Recommendation on Somatic Acupuncture




Recommendation on Scalp Acupuncture




Recommendation on Ear Acupuncture^[1]

 Lung Point (Required key point, alternate ear) Improves the function of lungs.
 Zero Point (add as needed, alternate ear) help relieving pain, and certain allergic reaction.
 Shenmen (add as needed, alternate ear) have tranquilizing effect, for anxiety, insomnia, HBP.


Procedure please follow the ear acupuncture operation routine, and the requirement of different auricular acupoints.






Lung point

Location: In the inferior hemiconcha and lies in the most
concave part of the inner
foothill of the antihelix.
Indication: Coughing, asthma,
hoarse voice, acne, urticarial.



Zero Point

Location: Point Zero is located on a notch near the base of the helix root where it begins to rise. **Indication:** It is one of the most recognized auricular points and is used in the treatment of many ailments. Zero Point is known to have a powerful influence in treating various conditions including pain, anxiety, drug addiction, and inflammation.




Shen men

Location: lies in the triangular
fossa and sits on the origin of
the superior and inferior
branches of the antihelix.
Indication: anxiety, stress, HBP,
depression, insomnia, pain,
addiction.

Recommendation on Somatic Acupuncture^[2,3]


Key point: ST36 (zu san li), required, help restore the balance of immune system ^[4,5] and Improve gastrointestinal symptoms.
Low fever: add LI11 (qu chi)
Chest tightness: add PC6 (nei guan)
Cough or difficult breathing: add LU6 (kong zui)
Lower GI complaints, e.g. diarrhea, constipation, bloating: add ST25 (tian shu) ^[6]
High fever: add DU14 (da zhui) ^[7,8]

Procedure Please follow the acupuncture operation routine and the acupuncture requirements of different acupoints.



Required Key point:ST36 (Zu san li)


- Foot Yangming Meridian Acupuncture Points,
- Location: On the lateral of the lower
 leg, 3 cun below ST 35, on the line
 joining ST 35 and ST 41.
 Indications: Gastrointestinal disease,
 e. g. abdominal pain, bloating,
 diarrhea, and constipation, and help
 restore the balance of immune system.
 Application: Perpendicular insertion
 0.7-1.5 inch.




Low fever: add LI11(Qu chi)

- Hand Taiyin Meridian Acupuncture Points.
- **Location:** At the midpoint of the line joining the lateral humeral epicondyle and the center of the cubital crease.
- Indications: Fever and febrile disease, allergy, sore throat, eye pain with redness, rubella(measles), upper limb motor impairment. Application: Perpendicular insertion 1.0-1.5 inch.





Low fever add LI11 (qu chi)



Chest tightness:add PC6 (nei guan)


Hand Jueyin Meridian Acupuncture Points. **Location:** On the palmar aspect of the forearm, 2 cun above the transverse crease of the wrist, between the tendons of m.palmaris longus and m.flexor carpi radialis. Indications:Cardiac pain, palpitations, nausea and vomiting ^[9], stomachache, elbow and arm soreness. **Application:** Perpendicular insertion 0.5-0.8 inch.

Cough or difficult breathing: add LU6 (kong zui)

Hand Taiyin Meridian Acupuncture Points


Location: At the front of the forearm, on the line joining the lateral end of the wrist and the center of the cubital crease, 7 cun above the transverse crease of the wrist.

Indications: Hemoptysis (coughing blood), epistaxis(nosebleed), cough and wheezing, sore throat.
Application: Perpendicular insertion 0.5-1.0 inch.



Celestial Pivot: add ST25 (tian shu)


Foot Yangming Meridian Acupuncture Points. Location: On the abdomen, horizontal to the umbilicus, 2 cun lateral to the anterior midline. Indications: Abdominal pain, constipation, diarrhea, irregular menstruation, edema. Application: Perpendicular insertion 0.7-1.2 inch.



High fever: add DU14 (da zhui)

Du Meridian Acupuncture Points. Location: On the posterior midline, in the depression just below C7 spinous process.

Indications: Malaria, fever and febrile diseases, cough and wheezing, spine and back stiffness and pain.
Application: Oblique insertion upward 0.5-1.0 inch.



High fever add DU14 (da zhui)

Recommendation on Scalp Acupuncture

Foot Motor and Sensory Area


2 parallel needles, help improve the immune system, enhance physical strength, work like a tranquilizer, relieve pain, and calm anxiety.

Thoracic Cavity Area

One needle, for coughing, difficult breathing, and chest tightness.

Procedure


Please follow the scalp acupuncture operation routine.



Foot Motor and Sensory Area

Location:1 cm apart from each sideof the midpoint of the anterior andposterior midline, leading backward3 cm horizontal line.

Indications: Contralateral lower limb pain, numbness, paralysis, acute lumbar sprain, cortical polyuria, nocturnal urination, uterine prolapse, etc.



Thoracic Cavity Area

Location:

Between the gastric area and the anterior and posterior midline, with a straight line of 2 cm above and below the hairline.

Indications:

Coughing, difficult breathing, and chest tightness, and bronchial asthma, etc.

Thoracic Cavity Area

References

- [1] Zhi, W. Iris, et al. "Acupuncture for bortezomib-induced peripheral neuropathy: not just for pain." Integrative cancer therapies 17.4 (2018): 1079-1086. <u>https://doi.org/10.1177%2F1534735418788667</u>
- [2] Liu, Wei-Hong et al. "Understanding of guidance for acupuncture and moxibustion interventions on COVID-19 (Second edition) issued by CAAM." World Journal of Acupuncture-Moxibustion, 10.1016/j.wjam.2020.03.005. 17 Mar. 2020. <u>https://doi.org/10.1016/j.wjam.2020.03.005</u>
- [3] Deare JC, Zheng Z, Xue CCL, Liu JP, Shang J, Scott SW, Littlejohn G. Acupuncture for treating fibromyalgia. Cochrane Database of Systematic Reviews 2013, Issue 5. Art. No.: CD007070. https://www.cochrane.org/CD007070/MUSKEL acupuncture-for-fibromyalgia
- [4] Jianli, C. "The effect of acupuncture on serum IgE level in patients with chronic urticaria." Journal of traditional Chinese medicine= Chung i tsa chih ying wen pan 26.3 (2006): 189-190. <u>https://europepmc.org/article/med/17078446</u>
- [5] Yang, Guang, et al. "Effects of electro-acupuncture at Zusanli, Guanyuan for sepsis patients and its mechanism through immune regulation." Chinese journal of integrative medicine 22.3 (2016): 219-224. <u>https://doi.org/10.1007/s11655-016-2462-9</u>
- [6] Diehl, D L. "Acupuncture for gastrointestinal and hepatobiliary disorders." Journal of alternative and complementary medicine (New York, N.Y.) vol. 5,1 (1999): 27-45. <u>https://doi.org/10.1089/acm.1999.5.27</u>
- [7] Xiao, Lei, et al. "Clinical observation on effects of acupuncture at Dazhui (GV 14) for abating fever of common cold." Zhongguo zhen jiu= Chinese acupuncture & moxibustion 27.3 (2007): 169-172. https://www.ncbi.nlm.nih.gov/pubmed/?term=PMID: 17432639
- [8] Kwon, SeungWon, et al. "Acupuncture therapy for fever induced by viral upper respiratory tract infection (URTI) in military medical service: a case series." Acupuncture in Medicine 32.6 (2014): 509-511. <u>https://doi.org/10.1136%2Facupmed-2014-010629</u>
- [9] Lee, Anna, Simon KC Chan, and Lawrence TY Fan. "Stimulation of the wrist acupuncture point PC6 for preventing postoperative nausea and vomiting." Cochrane database of systematic reviews 11 (2015). https://doi.org/10.1002/14651858.CD003281.pub4

Experts contributed in making the recommendations

Editedi

Jun Hu,Ph.D.LAc. Bing Zhu,Ph.D.M.D.(China) Wen Zhou Zhong,M.S.LAc. Linda Lingzhi Hao,Ph.D,D.O.M. Ping Xu,Prof.M.D.(China)